

AD-A075 404

METEOROLOGY INTERNATIONAL INC MONTEREY CALIF

F/G 4/2

MONTHLY MEAN AND STANDARD DEVIATION FIELDS OF SEA-LEVEL PRESSUR--ETC(U)

OCT 78 F G CATON, M J CUMING

N62271-78-M-1644

UNCLASSIFIED

MII-M-235


NL

| OF |

AD  
A075404



END  
DATE  
FILMED  
11-79  
DDC



MICROCOPY RESOLUTION TEST CHART  
NATIONAL BUREAU OF STANDARDS-1963-A



# LEVEL

17  
13

Technical Report

MII Project-M-235

October 1978

AD A 075404

DDC  
RECEIVED  
OCT 22 1978  
E

MONTHLY MEAN AND  
STANDARD DEVIATION FIELDS OF  
SEA-LEVEL PRESSURE AND SURFACE WIND  
FOR THE NORTHERN HEMISPHERE

Technical Report

by

Francis G. Caton  
Michael J. Cuming

Meteorology International Incorporated  
2600 Garden Road, Suite 145  
Monterey, California 93940

Prepared for

Commanding Officer  
Fleet Numerical Weather Central  
Monterey, California 93940

Contract No. N62271-78-M-1644

This document has been approved  
for public release and sale; its  
distribution is unlimited.

79 22 19 024

## **DISCLAIMER NOTICE**

**THIS DOCUMENT IS BEST QUALITY  
PRACTICABLE. THE COPY FURNISHED  
TO DDC CONTAINED A SIGNIFICANT  
NUMBER OF PAGES WHICH DO NOT  
REPRODUCE LEGIBLY.**



UNCLASSIFIED

SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)

REPORT DOCUMENTATION PAGE		READ INSTRUCTIONS BEFORE COMPLETING FORM
1. REPORT NUMBER Technical Report	2. GOVT ACCESSION NO.	3. RECIPIENT'S CATALOG NUMBER
4. TITLE (and Subtitle) MONTHLY MEAN AND STANDARD DEVIATION FIELDS OF SEA-LEVEL PRESSURE AND SURFACE WIND FOR THE NORTHERN HEMISPHERE		5. TYPE OF REPORT & PERIOD COVERED
7. AUTHOR(s) Francis G. Caton Michael J. Cuming		6. PERFORMING ORG. REPORT NUMBER M-235
9. PERFORMING ORGANIZATION NAME AND ADDRESS Meteorology International Incorporated 2600 Garden Road, Suite 145 Monterey, California 93940		8. CONTRACT OR GRANT NUMBER(s) N62271-78-M-1644
11. CONTROLLING OFFICE NAME AND ADDRESS Fleet Numerical Weather Central Monterey, California 93940		10. PROGRAM ELEMENT, PROJECT, TASK AREA & WORK UNIT NUMBERS
14. MONITORING AGENCY NAME & ADDRESS (if different from Controlling Office)		12. REPORT DATE October 1978
		13. NUMBER OF PAGES 18
		15. SECURITY CLASS. (of this report) UNCLASSIFIED
		15a. DECLASSIFICATION/DOWNGRADING SCHEDULE
16. DISTRIBUTION STATEMENT (of this Report) <div style="border: 1px solid black; padding: 5px; text-align: center;">This document has been approved for public release and sale; its distribution is unlimited.</div>		
17. DISTRIBUTION STATEMENT (of the abstract entered in Block 20, if different from Report)		
18. SUPPLEMENTARY NOTES		
19. KEY WORDS (Continue on reverse side if necessary and identify by block number) Sea-Level Pressure.      Northern Hemisphere Surface Wind      Marine History 1946-1977 Time-Averaged Mean Values      Climatology Standard Deviation      Analysis Data Sets      Fields by Information Blending		
20. ABSTRACT (Continue on reverse side if necessary and identify by block number) In March 1978 there became available a Northern Hemisphere history of analyzed sea-level pressure fields and diagnosed wind fields covering the period January 1946 through December 1977 at 6-hourly intervals. These fields have been used to generate Data Sets containing monthly-mean, and all-years monthly-mean, fields of sea-level pressure and surface wind components together with associated fields of standard deviation. This Report is designed to be suitable for dissemination with data charts and/or		

UNCLASSIFIED

SECURITY CLASSIFICATION OF THIS PAGE(When Data Entered)

Continuation of Block 20. Abstract

Can → data tape fields and provides background information, format specifications and contents of the new Data Sets. All-years monthly-mean fields of sea-level pressure and surface wind (speed and direction) are presented in the Appendices.

Accession For	
NTIS GRA&I	<input checked="checked" type="checkbox"/>
DDC TAB	<input type="checkbox"/>
Unannounced	<input type="checkbox"/>
Justification	<i>for</i>
<i>from 50 km file</i>	
By	
Distribution/	
Availability Codes	
Dist.	Avail and/or special
<i>A</i>	<i>23</i>

UNCLASSIFIED

SECURITY CLASSIFICATION OF THIS PAGE(When Data Entered)

## TABLE OF CONTENTS

	<u>Page</u>
1. INTRODUCTION . . . . .	1
2. BACKGROUND OF THE MARINE HISTORY OF SEA-LEVEL PRESSURE AND DIAGNOSED WIND FIELDS . . . . .	3
2.1 Data Sources . . . . .	3
2.2 The Analysis System . . . . .	5
2.3 The Wind Fields . . . . .	5
3. THE TIME-AVERAGED MEAN FIELDS OF SEA-LEVEL PRESSURE . . . . .	7
3.1 Source Fields . . . . .	7
3.2 The SLP Data Sets . . . . .	7
3.2.1 Data Set SLP1 . . . . .	7
3.2.2 Data Set SLP2 . . . . .	9
3.2.3 Data Set SLP3 . . . . .	10
4. THE TIME-AVERAGED MEAN FIELDS OF WIND COMPONENTS .	12
4.1 Source Fields . . . . .	12
4.2 The Wind Data Sets . . . . .	12
4.2.1 Data Set WUV1 . . . . .	12
4.2.2 Data Set WUV2 . . . . .	14
4.2.3 Data Set WUV3 . . . . .	16
5. REFERENCES . . . . .	18



TABLE OF CONTENTS (Continued)

APPENDIX A	DATA FORMAT SPECIFICATIONS AND PLOT ROUTINE
APPENDIX B	ALL-YEARS MONTHLY MEAN FIELDS OF SEA-LEVEL PRESSURE
APPENDIX C	ALL-YEARS MONTHLY MEAN FIELDS OF SURFACE WIND

## 1. INTRODUCTION

In March 1978, Meteorology International Incorporated completed a series of projects on behalf of FNWC which culminated, in part, in a Northern Hemisphere history of analyzed sea-level pressure fields<sup>1</sup> and diagnosed wind fields covering the period January 1946 through December 1977 in 6-hourly intervals--a total of some 47,000 fields for each of the two parameters. In effect these fields provide a time-sequence of sea-level pressure and diagnosed wind every 6 hours for a 32-year period at any point in the Northern Hemisphere. The fields<sup>2</sup> of sea-level pressure were produced by applying a consistent state-of-the-art analysis technique (the Fields by Information Blending methodology) to a very extensive data base (Tape Data Family-11 and Supplements, augmented by FNWC data sets). The wind fields were diagnosed from the fields of sea-level pressure.

Time-averaged mean fields of sea-level pressure and surface wind, together with associated fields of standard deviation, are required for many meteorological purposes ranging, for example, from studies of climatic trends to the evaluation of certain aspects of extended-range forecasting techniques. The best fields of mean sea-level pressure and wind hitherto available for such purposes have been compiled from sequences of real-time synoptic analyses. However in marine areas the real-time observational data contributing to these analyses was much less comprehensive than the data collections available for the reanalysis of Marine History and, in

---

<sup>1</sup>Known as the "Marine History" because of the emphasis placed on improving the quality of historical sea-level pressure analyses in marine areas.

<sup>2</sup>All fields referred to in this Report utilize the standard FNWC 63x63 analysis grid, polar stereographic projection.

addition, the analysis techniques employed changed with time as improvements were made and implemented. Thus one readily-apparent application of the Marine History is the determination of improved time-averaged mean fields of sea-level pressure and surface wind, together with associated fields of standard deviation.

These improved fields have been compiled from the Marine History and this Report, designed to be suitable for dissemination with data charts and/or data tape files, provides background information and format specifications<sup>3</sup> for the new data sets. Details of the Fields by Information Blending (FIB) methodology, the production of the Marine History, and the wind algorithm, are to be found in the References.

---

<sup>3</sup>Format specifications are given in Appendix A.



## 2. BACKGROUND OF THE MARINE HISTORY OF SEA-LEVEL PRESSURE AND DIAGNOSED WIND FIELDS

### 2.1 Data Sources

The FNWC copy of TDF-11, including Supplements, occupies over 550 large-reel magnetic tapes, the total number of reports being in excess of 50 million for the period 1854-1974. The reports in this FNWC "Marine Deck" are organized by Marsden square; within each Marsden square they are organized by year, and then by month. This organization is suitable for the compilation of climatological summaries but is inappropriate for synoptic analyses.

To establish a synoptic marine data base necessary for a reanalysis of Marine History, the FNWC Marine Deck was processed to extract, synoptically organize, and compact the observed values of sea-level pressure, and wind direction and speed, for the period 1946-1972. Data for the years 1973 to 1975 required to complete the 30-year period 1946-1975 was to be provided by marine data obtained by FNWC during the course of conducting routine operational synoptic analyses of sea-level pressure. However, for reasons given in Reference [1], the procedure eventually adopted was to reanalyze to 1971 and then to "add on" the high quality FIB analyses then being produced by FNWC to cover the period 1972-1975. An additional two years was subsequently added to cover the 32-year period 1946-1977.

Any hemispheric analysis of sea-level pressure cannot be performed using marine observations only. To carry out the reanalysis land data was essential, particularly from island and coastal stations. However no ready-made collection of land reports was available which could make any significant contribution to the reanalysis and, because of the magnitude of the task, such a collection could not be undertaken.

The problem of acquiring the information content provided by land observations, without actually assembling a data base of individual reports, was resolved by the use of historical surface-pressure analyses--these fields include all relevant land, island and ship reports which were available at the time the analysis was performed. It was not known, however, which stations actually contributed reports to each synoptic analysis of the available history. How the problems of extracting the essential information from each historical field and effecting its assimilation into the reanalysis were resolved is described in Reference [1].

Another source of information for any particular analysis of a meteorological parameter is the preceding analysis; i.e., information that has been carried forward along the time axis. The steering field necessary to carry information along the time axis by kinematical extrapolation was provided by the  $SR_{500}$  field [2].

The input data for analysis at each map time thus consisted of:

- a. Ship reports from the synoptically organized Marine Deck.
- b. The history field for the current map time. (Missing history fields were generated by forward-and-backward extrapolation.)
- c. The field previously analyzed. (In order to maximize the yield and utilization of information, the reanalysis was designed to progress backward-in-time.)
- d. The steering field based on the geostrophic wind derived from the SR-component of the 500-mb height field nearest to the current map time.

## 2.2 The Analysis System

The analysis system utilized for the reanalysis of Marine History was an application of the Fields by Information Blending methodology [3]. The data sources described above were provided as input to the system, which generated 6-hourly fields of sea-level pressure on a 63x63 analysis grid for the Northern Hemisphere. These fields were stored on magnetic tape, initially in a high-density format (9-track tape; 1600 b.p.i.). Subsequently this format was converted to NEDN format<sup>1</sup> (7-track tape; 800 b.p.i.) this storage density allowing one year of Marine History to be stored on each tape.

## 2.3 The Wind Fields

Wind fields (speed and direction) were diagnosed for the FNWC 63x63 analysis grid from the pressure fields using the algorithm originally developed for use in the FNWC Singular Sea/Swell Model [4]. The wind fields were stored on magnetic tape in NEDN format.

Prior to production the wind algorithm was examined in detail by both FNWC and MII with a view to effecting improvements. A comparison was made between over 2.3 million wind speeds reported by ships with the corresponding diagnosed wind speeds; no significant statistical bias was uncovered.<sup>2</sup> Other independent investigators had previously established that the wind algorithm produces no systematic bias with regard to wind direction. It was concluded therefore that no modification to the formulation could be devised which gave a significant overall improvement.

---

<sup>1</sup> See Appendix A.

<sup>2</sup> On average, diagnosed wind speeds were found to be approximately 1.8% low when compared with observed wind speeds.



It should be noted that the wind algorithm computes a quasi-geostrophic wind from the pressure field. The algorithm has been tuned so that the computed values of wind are representative of observed surface winds in marine areas. For land areas the computed winds should be interpreted as quasi-geostrophic winds--they are not (necessarily) representative of observed surface winds over land.

In addition, the wind defined by the pressure gradient in tropical regions may not be representative of observed winds for two reasons:

- a. The geostrophic wind relationship becomes less meaningful in low latitudes.
- b. Tropical cyclones are generally below the resolution capabilities of the grid used for pressure analysis and hence will not be represented in the diagnosed wind field.

### 3. THE TIME-AVERAGED MEAN FIELDS OF SEA-LEVEL PRESSURE

#### 3.1 Source Fields

These consisted of the 46,752 analyzed fields of sea-level pressure for the Northern Hemisphere covering the period 00Z 01JAN46 to 18Z 31DEC77 in 6-hourly intervals, each field being represented by 63x63 grid points on a polar stereographic projection.

Three data sets of time-averaged mean fields and associated fields of standard deviation were compiled from the source fields.

#### 3.2 The SLP Data Sets

##### 3.2.1 Data Set SLP1

This contains 384 fields of sea-level pressure, each an average over one calendar month from JAN 46 to DEC 77, and ordered chronologically. Each field of SLP is accompanied by an associated field of standard deviation.

For a calendar month of 30 days, for example, 120 source fields (30 days x 4 analyses per day) contributed to the monthly mean value of sea-level pressure; these 120 fields were used to compute the associated field of standard deviation for that particular month.

Table 3.1 shows the organization of Data Set SLP1.

Table 3.1  
Organization of Data Set SLP1.

	JAN	FEB	MAR	----	NOV	DEC
1946	$\overline{\text{SLP}}_J, \sigma_J$	$\overline{\text{SLP}}_F, \sigma_F$				$\overline{\text{SLP}}_D, \sigma_D$
1947	$\overline{\text{SLP}}_J, \sigma_J$	$\overline{\text{SLP}}_F, \sigma_F$				$\overline{\text{SLP}}_D, \sigma_D$
1948 ⋮ 1976						
1977	$\overline{\text{SLP}}_J, \sigma_J$	$\overline{\text{SLP}}_F, \sigma_F$				$\overline{\text{SLP}}_D, \sigma_D$

( $\overline{\text{SLP}}_M$  denotes the monthly mean field of SLP for month M,  $\sigma_M$  being the associated field of standard deviation.)

The 384 fields of  $\overline{\text{SLP}}_M$  and the 384 fields of  $\sigma_M$ , arranged in pairs in accordance with Table 3.1, are stored on one large-reel magnetic tape in NEDN format.



### 3.2.2 Data Set SLP2

This contains the same 384 pairs of fields as Data Set SLP1 but ordered by like-month rather than chronologically.

Table 3.2  
Organization of Data Set SLP2.

	1946	1947	1948 ---- 1976	1977
JAN	$\overline{\text{SLP}}_J, \sigma_J$	$\overline{\text{SLP}}_J, \sigma_J$		$\overline{\text{SLP}}_J, \sigma_J$
FEB	$\overline{\text{SLP}}_F, \sigma_F$	$\overline{\text{SLP}}_F, \sigma_F$		$\overline{\text{SLP}}_F, \sigma_F$
MAR : NOV				
DEC	$\overline{\text{SLP}}_D, \sigma_D$	$\overline{\text{SLP}}_D, \sigma_D$		$\overline{\text{SLP}}_D, \sigma_D$

( $\overline{\text{SLP}}_M$  denotes the monthly mean field of SLP for month  $M$ ,  $\sigma_M$  being the associated field of standard deviation.)

The 384 fields of  $\overline{\text{SLP}}_M$  and the 384 fields of  $\sigma_M$ , arranged in pairs in accordance with Table 3.2, are stored on one large-reel magnetic tape in NEDN format.

### 3.2.3 Data Set SLP3

This contains 12 fields of sea-level pressure, one for each calendar month and averaged over all 32 years. (These correspond to the "standard" monthly climatologies to be found in climatological atlases.) Each field is accompanied by an associated field of standard deviation.

For a calendar month of 30 days, for example, 3840 source fields (32 years x 30 days x 4 analyses per day) contributed to the all-years monthly mean field of sea-level pressure,  $\overline{\text{SLP}}_{\text{M}}$ . However these monthly fields actually were computed from the 32 appropriate fields of  $\overline{\text{SLP}}_{\text{M}}$ . Similarly, the associated all-years monthly fields of standard deviation were computed from the 32 pairs of appropriate fields of  $\sigma_{\text{M}}$  and  $\overline{\text{SLP}}_{\text{M}}$  rather than (unnecessarily) from the contributing source fields:

$$\sigma_{\text{M}}^2 = \frac{1}{32} \left[ \sum_{32} \sigma_{\text{M}}^2 + \sum_{32} \left( \overline{\text{SLP}}_{\text{M}} - \overline{\text{SLP}}_{\text{M}} \right)^2 \right]$$

at each grid point. It should be noted that the standard deviation corresponding to a mean field is that of all source fields making up the mean field.

Table 3.3 shows the organization of Data Set SLP3.

Table 3.3  
Organization of Data Set SLP3.

1946 to 1977	
JAN	$\overline{\text{SLP}}_{\text{J}}, \sigma_{\text{J}}$
FEB	$\overline{\text{SLP}}_{\text{F}}, \sigma_{\text{F}}$
MAR ⋮ NOV	
DEC	$\overline{\text{SLP}}_{\text{D}}, \sigma_{\text{D}}$

$\overline{\text{SLP}}_{\text{M}}$  denotes the all-years monthly mean field of SLP for calendar month M,  $\sigma_{\text{M}}$  being the associated field of standard deviation.)

The 12 fields of  $\overline{\text{SLP}}_{\text{M}}$ <sup>1</sup> and the 12 fields of  $\sigma_{\text{M}}$ , arranged in pairs in accordance with Table 3.3, are stored on one reel of magnetic tape in NEDN format.

---

<sup>1</sup>The all-years monthly mean fields of SLP are shown in Appendix B.

#### 4. THE TIME-AVERAGED MEAN FIELDS OF WIND COMPONENTS

##### 4.1 Source Fields

These consisted of the diagnosed fields of wind speed and direction interpreted in terms of  $u$  and  $v$  components where, by FNWC convention,  $u$  is directed along the  $I$ -axis of the analysis grid ( $u$  is positive in the direction of increasing  $I$ ) and  $v$  is directed along the  $J$ -axis of the analysis grid ( $v$  is positive in the direction of increasing  $J$ ).

Three data sets of time-averaged mean fields of surface wind and associated fields of standard deviation were compiled from the source fields.

##### 4.2 The Wind Data Sets

###### 4.2.1 Data Set WUV1

This contains 384 fields of the  $u$ -component of diagnosed wind, and 384 fields of the  $v$ -component of diagnosed wind, each of the 768 fields being an average over one calendar month from JAN46 to DEC77 and ordered chronologically. Each field is accompanied by an associated field of standard deviation.

For a 30-day calendar month, for example, 120 fields of the  $u$ -component of wind contributed to the monthly mean field of  $u$  and 120 fields of the  $v$ -component of wind contributed to the monthly mean field of  $v$ . The appropriate set of 120 component-fields were utilized to compute the standard deviations of the  $u$  and  $v$  components of wind respectively for that particular month.

Table 4.1 shows the organization of Data Set WUV1.



Table 4.1

Organization of Data Set WUV1.

	JAN	FEB	MAR ---- NOV	DEC
1946	$\overline{u_J}, \sigma_{uJ}; \overline{v_J}, \sigma_{vJ}$	$\overline{u_F}, \sigma_{uF}; \overline{v_F}, \sigma_{vF}$		$\overline{u_D}, \sigma_{uD}; \overline{v_D}, \sigma_{vD}$
1947	$\overline{u_J}, \sigma_{uJ}; \overline{v_J}, \sigma_{vJ}$	$\overline{u_F}, \sigma_{uF}; \overline{v_F}, \sigma_{vF}$		$\overline{u_D}, \sigma_{uD}; \overline{v_D}, \sigma_{vD}$
1948 : : 1976				
1977	$\overline{u_J}, \sigma_{uJ}; \overline{v_J}, \sigma_{vJ}$	$\overline{u_F}, \sigma_{uF}; \overline{v_F}, \sigma_{vF}$		$\overline{u_D}, \sigma_{uD}; \overline{v_D}, \sigma_{vD}$

$\overline{u_M}, \overline{v_M}$  denote the monthly mean fields of  $u$  and  $v$  respectively for month  $M$ ,  $\sigma_{uM}$  and  $\sigma_{vM}$  being the associated fields of standard deviation.)

The fields of  $\overline{u_M}$ ,  $\sigma_{uM}$  and  $\overline{v_M}$ ,  $\sigma_{vM}$ , arranged in pairs in accordance with Table 4.1, are stored on one large-reel magnetic tape in NEDN format.

#### 4.2.2 Data Set WUV2

This contains the same 768 pairs of fields as Data Set WUV1 but ordered by like-month rather than chronologically. Table 4.2 shows the organization of Data Set WUV2. The fields of  $\overline{u_M}$ ,  $\sigma_{uM}$  and  $\overline{v_M}$ ,  $\sigma_{vM}$ , arranged in pairs in accordance with Table 4.2, are stored on one large-reel magnetic tape in NEDN format.



Table 4.2  
Organization of Data Set WUV2.

	1946	1947	1948 ----- 1976	1977
JAN	$\overline{u}_J, \sigma_{uJ} ; \overline{v}_J, \sigma_{vJ}$	$\overline{u}_J, \sigma_{uJ} ; \overline{v}_J, \sigma_{vJ}$		$\overline{u}_J, \sigma_{uJ} ; \overline{v}_J, \sigma_{vJ}$
FEB	$\overline{u}_F, \sigma_{uF} ; \overline{v}_F, \sigma_{vF}$	$\overline{u}_F, \sigma_{uF} ; \overline{v}_F, \sigma_{vF}$		$\overline{u}_F, \sigma_{uF} ; \overline{v}_F, \sigma_{vF}$
MAR ⋮ NOV				
DEC	$\overline{u}_D, \sigma_{uD} ; \overline{v}_D, \sigma_{vD}$	$\overline{u}_D, \sigma_{uD} ; \overline{v}_D, \sigma_{vD}$		$\overline{u}_D, \sigma_{uD} ; \overline{v}_D, \sigma_{uD}$

( $\overline{u}_M, \overline{v}_M$  denote the monthly mean fields of  $u$  and  $v$  respectively for month  $M$ ,  $\sigma_{uM}$  and  $\sigma_{vM}$  being the associated fields of standard deviation.)

#### 4.2.3 Data Set WUV3

This contains 12 fields of the u-component of surface wind, one for each calendar month and averaged over all 32 years, and 12 fields of the v-component of surface wind, one for each calendar month and averaged over all 32 years. Each field is accompanied by an associated field of standard deviation.

For a calendar month of 30 days, for example, 3840 source fields (32 years x 30 days x 4 analyses per day) contributed to the all-years monthly mean field of each wind component. However, as with sea-level pressure, these monthly fields actually were computed from the appropriate mean fields. Similarly, the associated all-years monthly fields of standard deviation were computed from the appropriate mean fields (see Section 3.2.3).

Table 4.3 shows the organization of Data Set WUV3.

Table 4.3  
Organization of Data Set WUV3.

1946 - 1977	
JAN	$\overline{u_J}, \sigma_{\overline{u_J}}; \overline{v_J}, \sigma_{\overline{v_J}}$
FEB	$\overline{u_F}, \sigma_{\overline{u_F}}; \overline{v_F}, \sigma_{\overline{v_F}}$
MAR : NOV	
DEC	$\overline{u_D}, \sigma_{\overline{u_D}}; \overline{v_D}, \sigma_{\overline{v_D}}$

( $\overline{u_M}, \overline{v_M}$  denote the all-years monthly mean fields of  $u$  and  $v$  respectively for calendar month  $M$ ,  $\sigma_{\overline{u_M}}$  and  $\sigma_{\overline{v_M}}$  being the associated fields of standard deviation.)

The fields of  $\overline{u_M}, \sigma_{\overline{u_M}}$  and  $\overline{v_M}, \sigma_{\overline{v_M}}$ , arranged in pairs in accordance with Table 4.3, are stored on one reel of magnetic tape in NEDN format. All-years monthly mean fields of surface wind (speed and direction), computed from fields of  $\overline{u_M}$  and  $\overline{v_M}$ , are given in Appendix C.

5. REFERENCES

- [1] Mendenhall, Bruce R., Manfred M. Holl and Michael J. Cuming, Development of a Marine History of Analyzed Sea-Level Pressure Fields and Diagnosed Wind Fields, Technical Report, Contract N00228-76-C-3273, Meteorology International Incorporated, Monterey, California, 36 pp., 1978.
  
- [2] Holl, M. M., Scale-and-Pattern Spectra and Decompositions, Technical Memorandum No. 3, Contract N228-(62271)60550, Meteorology International Incorporated, Monterey, California, 28 pp., 1963.
  
- [3] Holl, Manfred M., The Upper-Air Analysis Capability, FIB/UA Introducing Weighted Spreading, Final Report, Contract N00228-75-C-2374, Meteorology International Incorporated, Monterey, California, 32 pp. plus Appendix A, 1976.
  
- [4] Hubert, W. E. and B. R. Mendenhall\* (\*Meteorology International Incorporated), The FNWC Singular Sea/Swell Model, Technical Note 59, Fleet Numerical Weather Central, Monterey, California, 29 pp., 1970.



APPENDIX A  
DATA FORMAT SPECIFICATIONS  
AND  
PLOT ROUTINE

## A.1 INTRODUCTION

The purpose of this Appendix is to provide users of the FNWC Monthly Mean and Standard Deviation Fields of Sea-Level Pressure and Surface Wind components for the Northern Hemisphere with a description of the data set format. Users of the Appendix are assumed to be familiar with the associated Technical Report.

## A.2 DATA SET CHARACTERISTICS

### A.2.1 Organization

The sea-level pressure data set is organized in the following tape file structure where each field is a record of the indicated file and mean and standard deviation fields are paired as consecutive records. Each file is contained on a single reel of magnetic tape.

- |      |   |
|------|---|
| SLP1 | Individual monthly mean and standard deviation fields organized chronologically from 1946 to 1977 (768 fields). |
| SLP2 | Individual monthly mean and standard deviation fields organized by like months (768 fields).                    |
| SLP3 | All-years monthly mean and standard deviation fields (24 fields).   |

The surface wind data set is organized similarly except that the U- and V-component mean fields and associated standard deviation fields are ordered as consecutive pairs. Each file is contained on a single magnetic tape.

- |      |  |
|------|--|
| WUV1 | Individual monthly mean and standard deviation fields organized chronologically from 1946 to 1977 (1536 fields). |
|------|--|



- WUV2      Individual monthly mean and standard deviation fields  
            organized by like months (1536 fields).
- WUV3      All-years monthly mean and standard deviation fields  
            (48 fields).

#### A.2.2 Data Tape Attributes

The data of each file has been recorded on 7-track tape at a density of 800 bits per inch. The tapes are unlabeled. The data have been written in Naval Environmental Data Network (NEDN) format. Specifics of NEDN format are provided in Sections A.2.3 and A.2.4. A full description of the NEDN format is presented in the FNWC User's Guide. Each data tape file is terminated by an end-of-information tape mark.

#### A.2.3 Data Characteristics

##### A.2.3.1 Packing Format

Each record of the files contains a packed version of a full field of 3969 (63x63) grid-point values and an associated identification (IDENT). The packed data field and IDENT are represented in 1007 60-bit words. These data are automatically unpacked if standard FNWC utility routines are used (see Section A.2.4). The first 14 60-bit words are used as the field IDENT; the remaining words contain the data points packed 4 to 1 (15 bits for each point). The following table identifies the pertinent elements of the field IDENT.

<u>PARAMETER</u>	<u>WORD (60 bit)</u>	<u>BIT POSITIONS</u>
Year (External BCD)	1	0-11
Month (External BCD)	2	48-59
Parameter Name (External BCD)	2	6-23
Scale Factor	3	1-6
This value represents the number of fractional bits (integer scaling) relative to a 60-bit word, for fixed point data values contained within the field. SLP fields = 36 WND fields = 40		
Field Title (External BCD)	9	0-36
	10-13	0-60

#### A.2.3.2 Parameter Units

The values of the sea-level pressure parameters, when unpacked and scaled, are expressed in millibars (to the nearest 1/8 mb); the wind parameters are expressed in meters per second (to the nearest 1/128 m/sec).

#### A.2.3.3 Parameter Names and Date-Time-Group Coding

For sea-level pressure fields, parameter names PMM (pressure monthly mean) and PMS (pressure monthly standard deviation) have been used. For wind parameters, UMM, UMS and VMM, VMS have been used for the U- and V-components of wind respectively.

Date-time-groups have been represented as the first day of the month for all individual monthly fields. For long-term monthly fields, the year and day equal zero. In all cases the hour has been set to zero.

#### A.2.4 FNWC Supporting Subroutines

FNWC maintains program library subroutines which are specially designed to handle NEDN data fields. Procedures for using these routines are provided in the FNWC Subroutine Manual. The routines are identified as follows:

<u>Subroutines</u>	<u>Purpose</u>
TW	To write 6500 fields from central memory to tape or disk with standard NEDN ident (readable with routine TR). Will perform float to fix conversion if necessary and pack the data as requested. Any size field may be written.
TR	To read NEDN standard format fields (written with routine TW) from tape or disk into central memory. Will convert ident to 6500 format, unpack the data and float if requested. There are two modes: a standard search under subroutine control; or a sequential search under user control.
TI	To print a list of the identification of fields written in NEDN format by subroutine TW.

#### A.3 UTILITY PLOT ROUTINE

A utility plot routine is available at FNWC for plotting full northern hemisphere contour charts for any of the fields provided in the data sets. The routine operates in a conversational mode in that it prompts the user in describing the fields he desires to be plotted.

Procedures for utilizing the plot routine simply require access to a remote terminal, execution of a BEGIN procedure and provision of the requested information. Specific access procedures can be obtained from the FNWC Data Archives and Display Division.



APPENDIX B


ALL-YEARS MONTHLY MEAN FIELDS  
OF SEA-LEVEL PRESSURE

## APPENDIX B

Appendix B contains all-years monthly mean fields of sea-level pressure for the Northern Hemisphere. These fields have been compiled from 6-hourly analyses on a 63x63 analysis grid, polar stereographic projection, covering the 32-year period 1946-1977--a total of 46,752 analyses. All analyses have been produced utilizing a consistent state-of-the-art analysis technique (an application of the Fields by Information Blending methodology). Monthly mean fields of sea-level pressure for the same period also are available as well as all associated fields of standard deviation.

### CONTENTS:

Figure B1	--	Mean sea-level pressure for JANUARY	, 1946-1977
B2	--	Mean sea-level pressure for FEBRUARY	, 1946-1977
B3	--	Mean sea-level pressure for MARCH	, 1946-1977
B4	--	Mean sea-level pressure for APRIL	, 1946-1977
B5	--	Mean sea-level pressure for MAY	, 1946-1977
B6	--	Mean sea-level pressure for JUNE	, 1946-1977
B7	--	Mean sea-level pressure for JULY	, 1946-1977
B8	--	Mean sea-level pressure for AUGUST	, 1946-1977
B9	--	Mean sea-level pressure for SEPTEMBER	, 1946-1977
B10	--	Mean sea-level pressure for OCTOBER	, 1946-1977
B11	--	Mean sea-level pressure for NOVEMBER	, 1946-1977
B12	--	Mean sea-level pressure for DECEMBER	, 1946-1977




Z 00 JAN 00 PMM

MARINE SLP MONTHLY MEANS-FNWC MII

Fig. B1

FLEET NUMERICAL WEATHER CENTRAL






12 00 FEB 00 PMM

MARINE SLP MONTHLY MEANS-FNWC MII

Fig. B2

FLEET METEOROL. WEATHER CENTRAL






Z 00 MAR 00 PMM  
Fig. B3

MARINE SLP MONTHLY MEANS-FNWC MII

FLEET NUMERICAL WEATHER CENTRAL



Z 00 APR 00 PMM

MARINE SLP MONTHLY MEANS-FNWC MII

Fig. B4

FLEET NUMERICAL WEATHER CENTRAL U.S.


Z 00 MAY 00 PMM

Fig. B5


MARINE SLP MONTHLY MEANS-FNWC MII

FLEET METEOROLOGICAL CENTER











Z 00 JUL 00 PMM

MARINE SLP MONTHLY MEANS-FNWC MII

FLEET NUMERICAL WEATHER CENTRAL

Fig. B7







MARINE SLP MONTHLY MEANS-FNWC MII

FLEET SUPERIOR WEATHER CENTER










12 00 NOV 00 PMM

MARINE SLP MONTHLY MEANS-FNWC MII

Fig. B11

FLEET SUPERIOR WEATHER CENTRAL



12-00 DEC 00 PMM  
Fig. B12

MARINE SLP MONTHLY MEANS-FNWC MII

FLEET NUMERICAL WEATHER CENTRAL

APPENDIX C

ALL-YEARS MONTHLY MEAN FIELDS  
OF SURFACE WIND



## APPENDIX C

Appendix C contains all-years monthly mean fields of surface wind (speed and direction) for the Northern Hemisphere computed from corresponding fields of the u and v components of wind. The wind-component fields were compiled from winds diagnosed from 6-hourly analyses of sea-level pressure on a 63x63 analysis grid, polar stereographic projection, covering the 32-year period 1946-1977--a total of 46,752 analyses. The sea-level pressure analyses were produced using a consistent state-of-the-art analysis technique (an application of the Fields by Information Blending methodology). The algorithm used to diagnose wind from sea-level pressure was tuned to provide a (quasi-geostrophic) wind representative of observed surface winds over marine areas. Over land the wind fields should be interpreted as quasi-geostrophic winds--they are not (necessarily) representative of observed surface winds for land areas. In addition, for reasons explained in the text, these winds are not representative in low latitudes (i.e., tropical regions).

The all-years monthly-mean surface wind vectors shown in Figs. C1 through C12 are represented by "arrows", the direction of the arrow being that of the wind. Wind speeds, to the nearest 5 knots, are shown by barbs with a long barb representing 10 knots and a short barb representing 5 knots.


Monthly mean fields of the u and v wind components for the same period also are available as well as all fields of standard deviation associated with wind component fields.

### CONTENTS:

- Figure C1 -- Vector mean wind for JANUARY , 1946-1977
- C2 -- Vector mean wind for FEBRUARY , 1946-1977
- C3 -- Vector mean wind for MARCH , 1946-1977




C4 -- Vector mean wind for APRIL , 1946-1977  
C5 -- Vector mean wind for MAY , 1946-1977  
C6 -- Vector mean wind for JUNE , 1946-1977  
C7 -- Vector mean wind for JULY , 1946-1977  
C8 -- Vector mean wind for AUGUST , 1946-1977  
C9 -- Vector mean wind for SEPTEMBER, 1946-1977  
C10 -- Vector mean wind for OCTOBER , 1946-1977  
C11 -- Vector mean wind for NOVEMBER , 1946-1977  
C12 -- Vector mean wind for DECEMBER , 1946-1977



00 JAN 00 MARINE WIND MONTHLY MEANS-FNWC MII 10/78

Fig. C1

FLEET NUMERICAL WEATHER CENTRAL - U.S. NAVY




00 FEB 00 MARINE WIND MONTHLY MEANS-FNWC MII 10/78

Fig. C2

FLEET NUMERICAL WEATHER CENTRAL, U.S. NAVY 11






10 MAR 00 MARINE WIND MONTHLY MEANS-FNWC MII 10/78

Fig. C3

FLEET NUMERICAL WEATHER CENTRAL, U.S. NAVY 19.27






10 APR 00 MARINE WIND MONTHLY MEANS-FNWC MII 10/78

Fig. C4

FLEET NUMERICAL WEATHER CENTRAL, U.S. NAVY 19.23




10 MAY 00 MARINE WIND MONTHLY MEANS-FNWC MII 10/78

Fig. C5

FLEET NUMERICAL WEATHER CENTRAL, U.S. NAVY 19.24.






30 JUN 00 MARINE WIND MONTHLY MEANS-FNWC MII 10/78

Fig. C6

FLEET NUMERICAL WEATHER CENTRAL, U.S. NAVY 19.24




00 JUL 00 MARINE WIND MONTHLY MEANS-FNWC MII 10/78

Fig. C7

FLEET NUMERICAL WEATHER CENTRAL, U.S. NAVY 19






30 AUG 00 MARINE WIND MONTHLY MEANS-FNWC MII 10/78

Fig. C8


FLEET NUMERICAL WEATHER CENTRAL U.S. NAVY 19



00 SEP 00 MARINE WIND MONTHLY MEANS-FNWC MII 10/78

Fig. C9

FLEET NUMERICAL WEATHER CENTRAL, U.S. NAVY 19.




00 OCT 00 MARINE WIND MONTHLY MEANS-FNWC MII 10/78

Fig. C10

FLEET NUMERICAL WEATHER CENTRAL U.S. NAVY 11






00 NOV 00 MARINE WIND MONTHLY MEANS-FNWC MII 10/78

Fig. C11

FLEET NUMERICAL WEATHER CENTRAL, U.S. NAVY 19





00 DEC 00 MARINE WIND MONTHLY MEANS-FNWC MII 10/78

Fig. C12

FLEET METEOROLOGICAL WEATHER CENTRAL, U.S. NAVY